

$2p$	M^2	A	$x_1 - x_2$	X	Mittel X	Beobachtetes τ	$\Delta\tau = \tau_2 - \tau_1$	$\Delta M = M_2^2 - M_1^2$
170+ 500	$10^{12} \times 3.757$	8.9 9.1	5.3 5.0	5.9 6.2	6.05	0.00115°		
170+ 750	$10^{12} \times 6.918$	15.2 16.0 15.4 15.7	7.4 10.0 8.3 9.0	11.0 10.3 10.3 10.6	10.55	0.00201°	0.00086°	$10^{12} \times 3.161$
170+1000	$10^{12} \times 10.945$	23.3 25.0 24.0 24.1	13.2 15.3 12.7 13.8	15.7 16.2 16.8 16.2	16.2	0.00309°	0.00108°	$10^{12} \times 4.027$
170+1250	$10^{12} \times 15.595$	31.8 33.7	16.8 18.0	22.2 23.4	22.8	0.00435°	0.00126°	$10^{12} \times 4.650$
170+1500	$10^{12} \times 20.980$	39.1 42.4 39.6 42.7	20.6 25.4 21.6 27.6	27.3 27.9 27.2 26.9	27.3	0.00520°	0.00085°	$10^{12} \times 5.385$
170+1750	$10^{12} \times 26.983$	48.2 52.5 49.0 50.7	26.6 32.8 27.8 31.6	33.0 33.7 33.1 32.7	33.1	0.00631°	0.00111°	$10^{12} \times 6.003$